

FEDERAL REPUBLIC OF GERMANY

Certificate of Priority for Filing of a Patent Application

Filing number: 100 39 719.0

Filing date: 14th August 2000

Applicant/Proprietor: Agfa-Gevaert Aktiengesellschaft,
Leverkusen/Germany

Title: A bleach-fixing agent bath
concentrate

Priority: 27.05.2000 DE 100 26 456.5 ---
03.08.2000 DE 100 37 765.3

IPC: G 03 C 5/38

The attached papers are a true and accurate reproduction of
the original documents for this patent application.

Munich, 7th November 2002

On behalf of the President
of the German Patent and Trade
Mark Office

(signature)

Hoiß

A bleach-fixing agent bath concentrate

This invention relates to a one-part bleach-fixing agent bath concentrate (BX concentrate) with which bleach-fixing agent baths can be made up or regenerated,
5 and also relates to a bleach-fixing agent bath.

BX baths are used in colour photographic processing in order to oxidise the metallic silver formed by development into a soluble form thereof (bleaching) and in order to dissolve it in this form, together with undeveloped silver halide, from the material
10 by complexation (fixing). For these purposes, BX baths contain a series of necessary chemicals, namely an iron(III) complex salt as an oxidant, a thiosulfate as a fixing agent, and a sulfite, a disulfite or a sulfinic acid as a stabiliser for the thiosulfate. These chemicals exert an effect on each other, so that they cannot be held for an extended period in the same solution. For example, the iron(III) complex salt
15 oxidises the sulfite, the disulfite or the sulfinic acid. The thiosulfate is thereby no longer stabilised, so that it then decomposes.

For this reason, BX baths are produced as two or three parts which are not combined with each other until just before they are used. Concentrates which are required for
20 regeneration, i.e. for replenishing chemicals which have been consumed, are likewise formulated as two or three parts.

Multi-part formulation of the constituents of a BX tank bath or of a BX regenerator is disadvantageous, firstly because it is costly and uneconomic, and secondly
25 because it results, time after time, in errors of addition.

There is therefore a great need for the chemicals for BX baths to be formulated as one part, and in particular there is a need to provide a one-part BX concentrate which can be converted very easily, namely by dilution with water, into a ready-to-
30 use BX bath, or which can be used just as easily for regenerating a BX bath. Attempts to satisfy these needs have hitherto failed because of the aforementioned decomposition of the thiosulfate, and also due to insufficient solubility of the

thiosulfate, of the sulfite and of the iron(III) complex salt, particularly if the latter is iron(III)-EDTA.

5 Surprisingly, it has now been found that these disadvantages can be overcome if at least one compound from the series comprising a phosphate, polyphosphate or polyphosphonate, or a nitrate or bromide, is added to the BX concentrate containing the aforementioned constituents.

10 Complex salts of Fe(III) which are suitable for photographic bleach and bleach-fixing baths are known from numerous documents (e.g. EP 329 088, 584 665, 507 126, 556 782, 532 003, 750 226, 657 777, 599 620, 588 289, 723 194, 851 287, 840 168, 871 065, 567 126, 726 203 and US 5 670 305).

15 The preferred complexing agents for Fe(III) are: ethylenediaminetetraacetic acid (EDTA), propylenediaminetetraacetic acid (PDTA), β -alaninediacetic acid (ADA), diethylenetriaminepentaacetic acid (DTPA), methyliminodiacetic acid (MIDA), ethylenediamine monosuccinate (EDMS), methylglycinediacetic acid (MGDA), ethylenediamine disuccinate (EDDS), particularly (S,S)-EDDS, iminosuccinic acid, iminosuccinic acid-propionic acid, and 2-hydroxypropyliminodiacetic acid.

20

Mixtures of complexing agents can also be used.

25 Examples of suitable sulfites include ammonium sulfite, ammonium hydrogen sulfite, sodium sulfite, sodium disulfite, sodium hydrogen sulfite, potassium sulfite, potassium disulfite and potassium hydrogen sulfite. Examples of suitable sulfinic acids include hydroxymethanesulfinic acid, formamidinesulfinic acid, benzenesulfinic acid, p-toluenesulfinic acid, methanesulfinic acid, o-amidosulfinic acid and salts thereof.

30 Alkali salts and/or ammonium salts can be used as phosphates, e.g. ammonium dihydrogen phosphate, diammonium hydrogen phosphate, triammonium phosphate, potassium dihydrogen phosphate, dipotassium hydrogen phosphate, tripotassium

phosphate, sodium dihydrogen phosphate, disodium hydrogen phosphate, and trisodium phosphate.

5 Examples of polyphosphates and polyphosphonates which can be used include sodium hexametaphosphate, sodium tetrphosphate, hydroxyethanediphosphonic acid, N(-2-carboxyethyl)-1-aminoethane-1,1-diphosphonic acid, N,N-bis-(carboxymethylene)-1-aminoethane-1,1-diphosphonic acid, morpholinomethane-diphosphonic acid, nitrilotrismethylene-phosphonic acid, ethylenediamine-tetramethylene phosphonic acid, hexamethylenediaminetetramethylene phosphonic
10 acid, 2-phosponobutane-1,2,4-tricarboxylic acid, and 2-carboxyethane-phosphonic acid. Free polyphosphoric acids are also suitable.

Alkali and/or ammonium nitrates and bromides can be used as nitrates and bromides.

15 The phosphates, polyphosphates and polyphosphonates, nitrates and bromides are preferably added to the concentrate in an amount ranging from 0.01 to 2.5 mol/litre, particularly from 0.05 to 1 mol/litre.

20 Sodium, potassium and ammonium thiosulfates are particularly suitable as fixing agents.

Other constituents can include aminopolycarboxylic acid, rehalogenating agents, acids and alkalies for pH adjustment, bleaching accelerators, white couplers and
25 buffer substances (see Research Disclosure 37 038, February 1995, pages 107 to 109).

In particular, the pH ranges from 4 to 9.

30 In addition, other complexing agents can also be added, individually or in admixture.

Polycarboxylic acids: e.g. oxalic acid, malonic acid, glutaric acid, adipic acid, suberic acid, fumaric acid, maleic acid, itaconic acid;

5 (Poly)hydroxypolycarboxylic acids: e.g. citric acid, glycolic acid, lactic acid, malic acid, tartaric acid, galactaric acid.

10 The present invention further relates to a ready-to-use bleach-fixing agent bath of the type cited at the outset, which is characterised in that it additionally contains a phosphate, particularly in an amount from 0.01 to 0.6 mol/litre, and a polyphosphate, particularly in an amount from 0.5 to 50 mmol/litre.

The bleach-fixing agent bath can be produced from the concentrate according to the invention if the concentrate contains phosphate and/or polyphosphate.

15

Examples**Example 1**

5 1 litre of BX concentrate contains

	ammonium thiosulfate solution, 57% by weight	400 ml
	ammonium hydrogen sulfite solution, 66% by weight	80 ml
	NH ₄ Fe(III)EDTA solution, 48% by weight	330 ml
10	additives	see below
	pH	5.5

The pH is adjusted with NH₃ or H₂SO₄.

15 The following additions were made to the BX concentrates:

BX 1: no additives

BX 2: 40 g/l sodium acetate (0.49 mol/litre)

BX 3: 186 g/l trisodium phosphate dodecahydrate (0.49 mol/litre)

BX 4: 50 g/l sodium hexametaphosphate (0.082 mol/litre)

20 BX 5: 73 ml/l aminotris(methylene)-phosphonic acid, concentration 50% by weight (0.16 mol/litre)

Storage at 60°C	Sodium sulfite content [g/l]				
Duration of storage	BX 1	BX 2	BX 3	BX 4	BX 5
No storage	82.7	82.5	82.4	82.6	82.3
2 days	55.6	56.0	65.4	64.9	65.1
6 days	Precipitates of sulfur		54.0	54.2	53.8

25 The stability of the sulfite is considerably improved by the addition of phosphate, polyphosphate and polyphosphonate.

The BX concentrate according to the invention can be used without disadvantages instead of a conventional, two-part BX concentrate, for example in the standard AP 94 process for the bleach-fixing of exposed, developed colour paper based on chloride-rich silver halide emulsions.

5

Example 2

The following additions were made to a BX concentrate as in Example 1 (no additives):

10

BX 1: no additives

BX 2: 40 g/l sodium acetate (0.49 mol/litre)

BX 3: 48.5 g/l ammonium dihydrogen phosphate (0.49 mol/litre)

BX 4: 48 g/l ammonium bromide (0.49 mol/litre)

BX 5: 73 g/l ammonium nitrate (0.49 mol/litre)

15

BX concentrate	Formation of crystals after 5 days at -5°C
BX 1	crystals
BX 2	crystals
BX 3	no crystals
BX 4	no crystals
BX 5	no crystals

The addition of phosphate, bromide or nitrate prevents the formation of crystals in a one-part bleach-fixing agent bath concentrate, so that a stable concentrate which comprises contents of active ingredients which would otherwise not be possible can also be produced.

20

The BX concentrate according to the invention is particularly suitable for short processing times (CD and BX times ranging from 12 to 35 seconds) and for a colour developer (CD) which contains disulfoethylhydroxylamine (HADS) as an antioxidant.

25

Example 3

A ready-to-use bleach-fixing agent bath was produced from the following components:

5

ammonium thiosulfate solution, 57% by weight 90 ml

sodium sulfite 10 g

NH₄ Fe(III)EDTA solution, 48 % by weight 70 ml

potassium dihydrogen phosphate 20 g

10

sodium hexametaphosphate 5 g

water to make up to 1 litre

Adjust pH to 6.5 with ammonia or phosphoric acid

15

This bleach-fixing agent bath is distinguished by the improved stability of the sulfite. It can be produced from a concentrate according to the invention.

Claims

1. A one-part photographic bleach-fixing agent bath concentrate containing an iron(III) complex salt, a thiosulfate and a sulfite, a disulfite or a sulfinic acid,
5 characterised in that it additionally contains at least one compound from the series comprising a phosphate, polyphosphate or polyphosphonate, or a nitrate or bromide.
2. A one-part bleach-fixing agent bath concentrate according to claim 1,
10 characterised in that its content of thiosulfate is 0.5 to 25 mol/litre, its content of sulfite is 0.2 to 2 mol/litre and its content of Fe(III) complex salt is 0.1 to 1 mol/litre.
3. A one-part bleach-fixing agent bath concentrate according to claims 1 or 2,
15 characterised in that its pH is 4 to 9.
4. A one-part bleach-fixing agent bath concentrate according to claims 1 or 2, characterised in that its pH is 5 to 6.5.
- 20 5. A one-part bleach-fixing agent bath concentrate according to any of claims 1 to 4, characterised in that the amount of phosphate, polyphosphate, polyphosphonate, nitrate or bromide is 0.01 to 2.5 mol/litre.
6. A one-part bleach-fixing agent bath concentrate according to any of claims 1
25 to 5, characterised in that it additionally contains one or more complexing agents.
7. A one-part photographic bleach-fixing agent bath concentrate containing an iron(III) complex salt, a thiosulfate, and a sulfite or a sulfinic acid,
30 characterised in that it additionally contains a phosphate, polyphosphate or polyphosphonate.

8. A ready-to-use bleach-fixing agent bath containing an iron(III) complex salt, a thiosulfate, and a sulfite, a disulfite or a sulfinic acid, characterised in that it additionally contains a phosphate and a polyphosphate.
- 5 9. A ready-to-use bleach-fixing agent bath according to claim 8, characterised in that the phosphate is contained in an amount from 0.01 to 0.6 mol and the polyphosphate is contained in an amount from 0.5 to 50 mmol per litre.
- 10 10. A ready-to-use bleach-fixing agent bath according to claims 8 and 9, characterised in that it is produced from a concentrate according to any of claims 1 to 7.

A bleach-fixing agent bath concentrate

Abstract

A one-part photographic bleach-fixing agent bath concentrate containing an iron(III) complex salt, a thiosulfate and a sulfite, a disulfite or a sulfinic acid, remains stable if a phosphate, polyphosphate or polyphosphonate, or a nitrate or bromide is added thereto.



**Prioritätsbescheinigung über die Einreichung
einer Patentanmeldung**

Aktenzeichen: 100 39 719.0

Anmeldetag: 14. August 2000

Anmelder/Inhaber: Agfa-Gevaert Aktiengesellschaft,
Leverkusen/DE

Bezeichnung: Bleichfixierbadkonzentrat

Priorität: 27.05.2000 DE 100 26 456.5
03.08.2000 DE 100 37 765.3

IPC: G 03 C 5/38

Die angehefteten Stücke sind eine richtige und genaue Wiedergabe der ursprünglichen Unterlagen dieser Patentanmeldung.

München, den 07. November 2002
Deutsches Patent- und Markenamt
Der Präsident
Im Auftrag

Hoß

Bleichfixierbadkonzentrat

Die Erfindung betrifft ein einteiliges Bleichfixierbadkonzentrat (BX-Konzentrat), mit dem Bleichfixierbäder angesetzt bzw. regeneriert werden können, sowie ein Bleichfixierbad.

BX-Bäder werden im farbfotografischen Verarbeitungsprozess eingesetzt, um das durch die Entwicklung entstandene metallische Silber zu einer löslichen Form zu oxidieren (Bleichung) und in dieser Form zusammen mit nicht entwickeltem Silberhalogenid durch Komplexbildung aus dem Material zu lösen (Fixierung). BX-Bäder enthalten für diese Aufgaben eine Reihe notwendiger Chemikalien, nämlich ein Eisen (III)-komplexsalz als Oxidationsmittel, ein Thiosulfat als Fixiermittel und ein Sulfid, Disulfid oder eine Sulfinsäure als Stabilisator für das Thiosulfat, die sich gegenseitig beeinflussen, so dass sie nicht längere Zeit in der gleichen Lösung gehalten werden können. Zum Beispiel oxidiert das Eisen(III)-komplexsalz das Sulfid, Disulfid oder die Sulfinsäure. Dadurch entfällt die Stabilisierung des Thiosulfates, das sich dann zersetzt.

Aus diesem Grund werden BX-Bäder zwei- oder dreiteilig konfektioniert, wobei die Teile erst unmittelbar vor dem Gebrauch vereinigt werden. Ebenso werden Konzentrate, die zum Regenerieren, d.h. zum Nachdosieren verbrauchter Chemikalien benötigt werden, zwei- oder dreiteilig konfektioniert.

Die mehrteilige Konfektionierung der Bestandteile einer BX-Tankfüllung oder eines BX-Regenerators ist nachteilig, weil sie einerseits aufwendig und unökonomisch ist, andererseits aber auch immer wieder zu Dosierungsfehlern führt.

Es besteht daher ein großer Bedarf, die Chemikalien für BX-Bäder einteilig zu konfektionieren und insbesondere ein einteiliges BX-Konzentrat bereitzustellen, das in einfachster Weise durch Verdünnen mit Wasser zum gebrauchsfertigen BX-Bad

umgesetzt oder ebenso einfach zum Regenerieren eines BX-Bad eingesetzt wird. Das scheiterte bisher an der vorstehend genannten Zersetzung des Thiosulfats, aber auch an der nicht ausreichenden Löslichkeit des Thiosulfates, des Sulfits und des Eisen(III)-komplexsalzes, insbesondere, wenn dieses Eisen(III)-EDTA ist.

Es wurde nun überraschend gefunden, dass diese Nachteile überwunden werden können, wenn man dem BX-Konzentrat mit den zuvor genannten Bestandteilen wenigstens eine Verbindung aus der Reihe Phosphat, Polyphosphat oder Polyphosphonat, Nitrat oder Bromid zusetzt.

Fe(III)-komplexsalze, die sich für fotografische Bleich- und Bleichfixierbäder eignen, sind aus einer Vielzahl von Dokumenten bekannt (z.B. EP 329 088, 584 665, 507 126, 556 782, 532 003, 750 226, 657 777, 599 620, 588 289, 723 194, 851 287, 840 168, 871 065, 567 126, 726 203 und US 5 670 305).

Bevorzugte Komplexbildner für Fe(III) sind: Ethylendiamintetraessigsäure (EDTA), Propylendiamintetraessigsäure (PDTA), β -Alanindiessigsäure (ADA), Diethyltri-aminpentaessigsäure (DTPA), Methyliminodiessigsäure (MIDA), Ethylendiaminmonosuccinat (EDMS), Methylglycindiessigsäure (MGDA), Ethylendiamindisuccinat (EDDS), speziell (S,S)-EDDS, Iminobernsteinsäure, Iminobernsteinsäurepropionsäure, 2-Hydroxypropyliminodiessigsäure.

Es sind auch Gemische von Komplexbildnern einsetzbar.

Als Sulfit eignen sich z.B. Ammoniumsulfit, Ammoniumhydrogensulfit, Natriumsulfit, Natriumdisulfit, Natriumhydrogensulfit, Kaliumsulfit, Kaliumdisulfit, Kaliumhydrogensulfit. Als Sulfinsäuren eignen sich z.B. Hydroxymethansulfinsäure, Formamidinsulfinsäure, Benzolsulfinsäure, p-Toluolsulfinsäure, Methansulfinsäure, o-Amidosulfinsäure und deren Salze.

Als Phosphate können die Alkalisalze und/oder Ammoniumsalze eingesetzt werden, z.B. Ammoniumdihydrogenphosphat, di-Ammoniumhydrogenphosphat, tri-Ammoniumphosphat, Kaliumdihydrogenphosphat, di-Kaliumhydrogenphosphat, tri-Kaliumphosphat, Natriumdihydrogenphosphat, di-Natriumhydrogenphosphat, tri-Natriumphosphat.

Als Polyphosphate und -Phosphonate können z.B. Natriumhexametaphosphat, Natriumtetraphosphat, Hydroxyethandiphosphonsäure, N(-2-carboxyethyl)-1-aminoethan-1,1-diphosphonsäure, N,N-Bis-(carboxymethylen)-1-aminoethan-1,1-diphosphonsäure, Morpholinomethandiphosphonsäure, Nitrilotrismethylenphosphonsäure, Ethylendiamintetramethylenphosphonsäure, Hexamethylendiamintetramethylenphosphonsäure, 2-Phosponobutan-1,2,4-tricarbonsäure, 2-Carboxyethanphosphonsäure eingesetzt werden. Geeignet sind auch freie Polyphosphorsäuren.

Als Nitrate und Bromide können Alkali- und/oder Ammoniumnitrate und -bromide eingesetzt werden.

Die Phosphate, Polyphosphate und Polyphosphonate, Nitrate und Bromide werden dem Konzentrat vorzugsweise in einer Menge von 0,01 bis 2,5 mol/l, insbesondere 0,05 bis 1 mol/l zugesetzt.

Als Fixiermittel eignen sich insbesondere Natrium-, Kalium- und Ammoniumthiosulfat.

Weitere Bestandteile können z.B. Aminopolycarbonsäure, Rehalogenierungsmittel, Säuren und Laugen zur pH-Einstellung, Bleichbeschleuniger, Weißkuppler und Puffersubstanzen sein (s. Research Disclosure 37 038, Februar 1995, Seiten 107 bis 109).

Der pH-Wert beträgt insbesondere 4 bis 9.

Zusätzlich können noch andere Komplexbildner einzeln oder im Gemisch zugesetzt werden:

Polycarbonsäuren: z.B. Oxalsäure, Malonsäure, Glutarsäure, Adipinsäure, Korksäure, Fumarsäure, Maleinsäure, Itaconsäure;

(Poly)Hydroxypolycarbonsäuren: z.B. Citronensäure, Glykolsäure, Milchsäure, Äpfelsäure, Weinsäure, Galactarsäure.

Ein weiterer Gegenstand der Erfindung ist ein gebrauchsfertiges Bleichfixierbad der eingangs genannten Art, das dadurch gekennzeichnet ist, dass es zusätzlich ein Phosphat, insbesondere 0,01 bis 0,6 mol/l, und ein Polyphosphat, insbesondere 0,5 bis 50 mmol/l, enthält.

Das Bleichfixierbad kann aus dem Konzentrat nach der Erfindung hergestellt werden, wenn das Konzentrat Phosphat und/oder Polyphosphat enthält.

Beispiele**Beispiel 1**

1 l BX-Konzentrat enthält

Ammoniumthiosulfatlösung, 57 Gew.-%	400 ml
Ammoniumhydrogensulfitlösung, 66 Gew.-%	80 ml
NH ₄ Fe(III)EDTA-Lösung, 48 Gew.-%	330 ml
Zusätze	siehe unten
pH-Wert	5,5

Die pH-Einstellung erfolgt mit NH₃ oder H₂SO₄.

Folgende Zusätze wurden den BX-Konzentraten zugegeben:

BX 1: ohne Zusätze

BX 2: 40 g/l Natriumacetat (0,49 mol/l)

BX 3: 186 g/l tri-Natriumphosphat-Dodecyhydrat (0,49 mol/l)

BX 4: 50 g/l Natriumhexametaphosphat (0,082 mol/l)

BX 5: 73 ml/l Aminotrismethylenphosphonsäure, 50 gew.-%ig (0,16 mol/l)

Lagerung bei 60°C	Natriumsulfitgehalt [g/l]				
Dauer der Lagerung	BX 1	BX 2	BX 3	BX 4	BX 5
Ungelagert	82,7	82,5	82,4	82,6	82,3
2 Tage	55,6	56,0	65,4	64,9	65,1
6 Tage	Schwefelausfällungen		54,0	54,2	53,8

Die Sulfitbeständigkeit wird durch den Zusatz von Phosphat, Polyphosphat und Polyphosphonat deutlich verbessert.

- 25 Das BX-Konzentrat gemäß der Erfindung kann ohne Nachteile anstelle eines herkömmlichen, zweiteiligen BX-Konzentrates eingesetzt werden, beispielsweise im

Standard-AP 94-Prozess zum Bleichfixieren von belichtetem und entwickeltem Colorpapier auf der Basis chloridreicher Silberhalogenidemulsionen.

Beispiel 2

Einem BX-Konzentrat nach Beispiel 1 (ohne Zusätze) werden folgende Zusätze zugegeben:

BX 1: ohne Zusätze

BX 2: 40 g/l Natriumacetat (0,49 mol/l)

BX 3: 48,5 g/l Ammoniumdihydrogenphosphat (0,49 mol/l)

BX 4: 48 g/l Ammoniumbromid (0,49 mol/l)

BX 5: 73 g/l Ammoniumnitrat (0,49 mol/l)

BX-Konzentrat	Bildung von Kristallen nach 5 Tagen bei -5°C
BX 1	Kristalle
BX 2	Kristalle
BX 3	keine Kristalle
BX 4	keine Kristalle
BX 5	keine Kristalle

Der Zusatz von Phosphat, Bromid oder Nitrat verhindert in einem einteiligen Bleichfixierbad-Konzentrat die Bildung von Kristallen, so dass auch mit sonst nicht möglichen Wirkstoffgehalten ein stabiles Konzentrat hergestellt werden kann.

Besonders geeignet ist das erfindungsgemäße BX-Konzentrat für kurze Verarbeitungszeiten (CD- und BX-Zeiten von 12 bis 35 s) und einen Farentwickler (CD), der Disulfoethylhydroxylamin (HADS) als Oxidationsschutzmittel enthält.

Beispiel 3

Es wird ein gebrauchsfertiges Bleichfixierbad aus folgenden Komponenten hergestellt:

Ammoniumthiosulfatlösung, 57 Gew.-%	90 ml
Natriumsulfit	10 g
NH ₄ Fe(III)EDTA-Lösung, 48 Gew.-%	70 ml
Kaliumdihydrogenphosphat	20 g
Natriumhexametaphosphat	5 g
Wasser aus 1 Liter	
pH-Wert mit Ammoniak oder Phosphorsäure auf 6,5 einstellen	

Das Bleichfixierbad zeichnet sich durch eine verbesserte Stabilität des Sulfits aus.

Die Herstellung kann aus einem erfindungsgemäßen Konzentrat erfolgen.

Patentansprüche

1. Einteiliges fotografisches Bleichfixierbadkonzentrat enthaltend ein Eisen(III)-komplexsalz, ein Thiosulfat und ein Sulfit, Disulfit oder eine Sulfinsäure, dadurch gekennzeichnet, dass es zusätzlich wenigstens eine Verbindung aus der Reihe Phosphat, Polyphosphat oder Polyphosphonat, Nitrat oder Bromid enthält.
2. Einteiliges Bleichfixierbadkonzentrat nach Anspruch 1, dadurch gekennzeichnet, dass sein Gehalt an Thiosulfat 0,5 bis 25 mol/l, sein Gehalt an Sulfit 0,2 bis 2 mol/l und sein Gehalt an Fe(III)-Komplexsalz 0,1 bis 1 mol/l beträgt.
3. Einteiliges Bleichfixierbadkonzentrat nach den Ansprüchen 1 oder 2, dadurch gekennzeichnet, dass sein pH-Wert 4 bis 9 beträgt.
4. Einteiliges Bleichfixierbadkonzentrat nach den Ansprüchen 1 oder 2, dadurch gekennzeichnet, dass sein pH-Wert 5 bis 6,5 beträgt.
5. Einteiliges Bleichfixierbadkonzentrat nach den Ansprüchen 1 bis 4, dadurch gekennzeichnet, dass die Menge an Phosphat, Polyphosphat, Polyphosphonat, Nitrat oder Bromid 0,01 bis 2,5 mol/l beträgt.
6. Einteiliges Bleichfixierbadkonzentrat nach den Ansprüchen 1 bis 5, dadurch gekennzeichnet, dass zusätzlich ein oder mehrere Komplexbildner enthalten sind.
7. Einteiliges fotografisches Bleichfixierbadkonzentrat enthaltend ein Eisen(III)-komplexsalz, ein Thiosulfat und ein Sulfit oder eine Sulfinsäure, dadurch gekennzeichnet, dass es zusätzlich ein Phosphat, Polyphosphat oder Polyphosphonat enthält.

8. Gebrauchsfertiges Bleichfixierbad enthaltend ein Eisen(III)-komplexsalz, ein Thiosulfat und ein Sulfit, Disulfit oder eine Sulfinsäure, dadurch gekennzeichnet, dass es zusätzlich ein Phosphat und ein Polyphosphat enthält.
9. Gebrauchsfertiges Bleichfixierbad nach Anspruch 8, dadurch gekennzeichnet, dass das Phosphat in einer Menge von 0,01 bis 0,6 mol und das Polyphosphat in einer Menge von 0,5 bis 50 mmol pro Liter enthalten ist.
10. Gebrauchsfertiges Bleichfixierbad nach Ansprüchen 8 und 9, dadurch gekennzeichnet, dass es aus einem Konzentrat nach den Ansprüchen 1 bis 7 hergestellt ist.

BleichfixierbadkonzentratZusammenfassung

Ein einteiliges fotografisches Bleichfixierbadkonzentrat enthaltend ein Eisen(III)-komplexsalz, ein Thiosulfat und ein Sulfit, Disulfit oder eine Sulfinsäure, bleibt stabil, wenn ihm ein Phosphat, Polyphosphat oder Polyphosphonat, ein Nitrat oder Bromid zugesetzt wird.


IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

DECLARATION

I, Paul David Churchill Clarke, BA. MITI., translator to Taylor & Meyer of 20 Kingsmead Road, London SW2 3JD, England, do solemnly and sincerely declare as follows:

1. That I am well acquainted with the English and German languages;
2. That the following is a true translation made by me into the English language of German Patent Application No. 100 37 765.3;
3. That all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardise the validity of the application or any patent issued thereon.

Signed, this 12 day of December 2002


Paul David Churchill Clarke
Harrogate, HG1 4QD, England